

**Ward, Liza**

---

**From:** Harwood, Valerie [vharwood@cas.usf.edu]  
**Sent:** Thursday, July 10, 2008 4:07 PM  
**To:** Ward, Liza; David Page  
**Subject:** FW: Manuscript submission (AEM01306-08 Version 1)

**Attachments:** PoultLitterQPCR\_MS\_FINAL.doc; AEMTMP-02130-08\_1[1].pdf



PoultLitterQPCRAEMTMP-02130-  
\_MS\_FINAL.doc .3\_1[1].pdf (140 .

Email forwarded as requested

Valerie J. (Jody) Harwood, Ph.D.  
Department of Biology, SCA 110  
University of South Florida  
4202 E. Fowler Ave.  
Tampa, FL 33620  
(813) 974-1524 - phone  
(813) 974-3263 - fax vharwood@cas.usf.edu  
4:07 PM

-----Original Message-----

**From:** Harwood, Valerie  
**Sent:** Wednesday, June 11, 2008 3:37 PM  
**To:** Jennifer Weidhaas (jweidhaas@northwind-inc.com); Tamzen MacBeth (tmacbeth@northwind-inc.com); Olsen Roger (olsenrl@cdm.com); David Page(new)  
**Subject:** FW: Manuscript submission (AEM01306-08 Version 1)

Manuscript submitted!!

Valerie J. (Jody) Harwood, Ph.D.  
Department of Biology, SCA 110  
University of South Florida  
4202 E. Fowler Ave.  
Tampa, FL 33620  
(813) 974-1524 - phone  
(813) 974-3263 - fax

-----Original Message-----

**From:** journalsrr@asmusa.org [mailto:journalsrr@asmusa.org]  
**Sent:** Wednesday, June 11, 2008 3:33 PM  
**To:** Harwood, Valerie  
**Subject:** Manuscript submission (AEM01306-08 Version 1)

Dr. Valerie Harwood  
University of South Florida  
Dept. of Biology  
4202 East Fowler Ave.  
Tampa, FL 33620-5550  
United States

Re: Identification and Validation of a Poultry Litter-Specific Biomarker and Development of a 16S rRNA Based Quantitative PCR Assay  
(AEM01306-08 Version 1)

Dear Dr. Harwood:

You have successfully submitted your manuscript via the Rapid Review system. The control number of your manuscript is AEM01306-08 Version 1. Take note of this number, and refer to it in any correspondence with the Journals Department or with the editor. You may log onto the Rapid Review system at any time to see the current status of your manuscript and the name of the editor handling it. The URL is <http://www.rapidreview.com/ASM2/author.html>, and your user name is vharwood. To find contact information for the editor handling your manuscript, go to the following URL: <http://www.asm.org/journals/editors.asp>

In submitting your manuscript to Applied and Environmental Microbiology (AEM), the author(s) guarantees that a manuscript with substantially the same content has not been submitted or published elsewhere and that all of the authors are aware of and agree to the submission.

By publishing in the journal, the authors agree that any DNAs, viruses, microbial strains, mutant animal strains, cell lines, antibodies, and similar materials newly described in the article are available from a national collection or will be made available in a timely fashion, at reasonable cost, and in limited quantities to members of the scientific community for noncommercial purposes. The authors guarantee that they have the authority to comply with this policy either directly or by means of material transfer agreements through the owner.

Similarly, the authors agree to make available computer programs, originating in the authors' laboratory, that are the only means of confirming the conclusions reported in the article but that are not available commercially. The program(s) and suitable documentation regarding its (their) use may be provided by any of the following means: (i) as a program transmitted via the Internet, (ii) as an Internet server-based tool, or (iii) as a compiled or assembled form on a suitable medium (e.g., magnetic or optical). It is expected that the material will be provided in a timely fashion and at reasonable cost to members of the scientific community for noncommercial purposes. The authors guarantee that they have the authority to comply with this policy either directly or by means of material transfer agreements through the owner.

If your manuscript is accepted for publication, a condition of acceptance is that you assign copyright to the American Society for Microbiology. A copyright transfer agreement is sent with each letter of acceptance after the manuscript has been scheduled for publication.

If your manuscript is accepted for publication in a 2008 issue, page charges (subject to change without notice) will be assessed at \$65 per printed page for the first eight pages and \$200 for each page in excess of eight for a corresponding author who is an ASM member or \$75 per printed page for the first eight pages and \$250 for each page in excess of eight for a nonmember corresponding author. A corresponding author who is not a member may join ASM to obtain the member rate. If the research was not supported, you may send a request for a waiver of page charges to the Director, Journals. For more details, including type of articles not charged, see the Instructions to Authors.

**IMPORTANT NOTICE:** For its primary-research journals, ASM posts online PDF versions of manuscripts that have been peer reviewed and accepted but not yet copyedited. This feature is called "AEM Accepts" and is accessible from the Journals website. The manuscripts are published online as soon as possible after acceptance, on a weekly basis, before the copyedited, typeset versions are published. They are posted "As Is" (i.e., as submitted by the authors at the modification stage), and corrections/changes are NOT accepted. Accordingly, there may be differences between the AEM Accepts version and the final, typeset version. The manuscripts remain listed on the AEM Accepts page until the final, typeset versions are posted, at which point they are removed from the AEM Accepts page and become available only through links from the final, typeset version. They are under subscription access control until 4 months after the typeset versions are posted, when access to all forms becomes free to everyone. Any supplemental material intended, and accepted, for publication is not posted until publication of the final, typeset article.

Thank you for submitting your manuscript for consideration.

Barbara Slinker  
Production Editor  
Applied and Environmental Microbiology (AEM)

**Identification and Validation of a Poultry Litter-Specific Biomarker and Development of a  
16S rRNA Based Quantitative PCR Assay**

Jennifer L. Weidhaas<sup>1</sup>, Tamzen W. Macbeth<sup>1</sup>, Roger L. Olsen<sup>2</sup>, Valerie J. Harwood<sup>3, \*</sup>

1. North Wind, Inc. 1425 Higham Street, Idaho Falls, Idaho, 83402

2. CDM, 555 17<sup>th</sup> St., Suite 1100, Denver, CO 80202

3. \* Department of Biology, University of South Florida, 4202 E. Fowler Ave., Tampa, Florida  
33620, Phone: 813-974-1524, Fax: 813-974-3263 email vharwood@cas.usf.edu

Running title: Brevibacterium marker for fecal source tracking of poultry

**ABSTRACT**

A poultry litter-specific biomarker was developed for microbial source tracking (MST) in environmental waters. 16S rRNA sequences that were present in fecal-contaminated turkey and chicken litter were identified by terminal restriction fragment length polymorphism (T-RFLP). Cloning and sequencing of potential targets from pools of *E. coli*, *Bacteroides* or total bacterial DNA yielded four sequences that were ubiquitous in poultry litter and also contained unique sequences for development of target-specific PCR primers. Primer sensitivity and specificity were tested by nested PCR against ten composite poultry litter samples and fecal samples from beef and dairy cattle, swine, ducks, geese, and human sewage. The sequence with greatest sensitivity (100%) and specificity (93.5%) has 98% identity to *Brevibacterium avium*, and was detected in all litter samples. It was detected at low level in only one goose and one duck sample. A quantitative PCR assay was developed and tested on litter, soil and water samples. Litter concentrations were  $2.2 \times 10^7$  -  $2.5 \times 10^9$  gene copies/g. The biomarker was present in a majority of soil and water samples collected in and near areas where litter was spread, reaching concentrations of  $2.9 \times 10^5$  gene copies·g<sup>-1</sup> in soil samples and  $5.5 \times 10^7$  gene copies·L<sup>-1</sup> in runoff from the edges of fields. The biomarker will contribute to quantifying the impact of fecal contamination by land-applied poultry litter in this watershed. Furthermore, it has potential for determining fecal source allocations for total maximum daily load (TMDL) programs and ambient water quality assessment, and may be useful in other geographic regions.

## INTRODUCTION

Excessive land application of poultry litter as a waste disposal mechanism has been linked to eutrophication of water bodies (28, 35, 39), the spread of pathogens (15, 19, 21), air and soil pollution with metals (11, 33) and groundwater contamination with nitrate (5). Despite these known effects, land application is still the typically practiced disposal method for poultry litter even though viable and economically favorable alternative disposal practices are available (7, 20).

Identification of the source of fecal pollution contaminating a watershed is of particular interest for protection of water resources and the safety of recreational waters. For example, TMDL assessments require identification of the source of contamination, which is also necessary for remediation of impaired waters(44). Current methods for detecting the presence of fecal pollution, which carries an increased risk of the presence of pathogenic microorganisms, involve the cultivation of fecal indicator organisms such as fecal coliforms in the family *Enterobacteriaceae* (Oklahoma Administrative Code, Title 785, Chapter 46). The U.S. EPA and many states recognize *Escherichia coli* and enterococci as indicators of freshwater recreational water quality (42).

Drawbacks to the use of indicator organisms which limit the ability of researchers to pinpoint sources of fecal contamination include the non-specificity of the fecal coliforms to one source (25, 43), variable survival rates of various indicator organisms (1) and the growth or extended persistence of these indicator organisms after release to the environment (12, 45). These drawbacks have lead to research into alternative methods for the assessment of human health risk

from microbial pathogens in recreational waters that do not include the culturing of fecal indicator organisms for identification and quantification of the source of fecal pollution (46).

A variety of microbial source tracking (MST) methods (for recent reviews see (17, 40, 47)) have been proposed as an alternative to cultivation of fecal coliforms. Some of these genotypic molecular based techniques have included library dependent methods (i.e., culture and isolate-based) such as ribotyping (10, 31) and repetitive element polymerase chain reaction (REP-PCR) (14). Library independent methods (i.e., detection of a genetic biomarker in extracted DNA) have also been developed using discovery techniques such as suspension arrays (8), subtractive hybridization (13, 26), and terminal restriction fragment length polymorphism (T-RFLP) (3), among others. Host marker specific targets have included *Enterococcus faecium* (37), *Bifidobacterium* and members of the *Bacteroidales* (3, 22, 38), among others. Relatively few microbial targets specific to poultry fecal material have been identified. To date *Enterococcus faecalis* (23), *E. coli* (10) and *Bacteriodes* (26) have been associated with poultry fecal material, but only the *Bacteroides* biomarker (26) was specifically associated with poultry and not other fecal sources. The objective of this research was to identify a poultry litter-specific biomarker, validate its specificity against other sources of fecal material from within and outside the watershed and develop a 16S rRNA based real-time PCR assay for quantifying the biomarker in environmental samples. This work was carried out as part of ongoing litigation in which the plaintiff is the Oklahoma Attorney General.

## **METHODS**

**Sample collection.** Litter samples were collected from ten separate facilities (poultry houses), nine chicken and one turkey facility. Litter samples were collected from 18 locations within each

poultry house through the entire depth of the litter. The subsamples (total volume of 4 to 5 gallons) from each house were composited, homogenized and split (riffle splitter) before placement into a sterile whirl pack (approximately 500 mL) and shipped on ice to the laboratory for analysis. Litter application areas in fields (soils) were sampled by collecting 20 subsamples on a predetermined grid pattern across a uniform subarea of one to ten acres in size. The zero to two inch sample from six inch soil cores were composited, disaggregated, sieved to 2 mm, ground, homogenized and split. Vegetation, feathers, and rocks were removed. The split soil samples (500 ml) were transported on ice to the laboratory. Nontarget fecal samples for specificity testing were collected as composites from groups of individuals (Table 3). Samples from beef cattle were collected from ten grazing fields, of which five were within the watershed and five were outside the watershed. Two independent duplicate samples were collected for each field, and each duplicate consisted of feces from ten scats. A total of 200 beef cattle scats were collected and composited into 20 samples. Duck and goose samples were collected in the same fashion, consisting of composites from ten individual scats, and independent duplicates were collected for each area. For ducks, three landing areas inside the watershed and two outside the watershed were sampled, while for geese, two landing areas inside and three landing areas outside the watershed were sampled. A total of 100 scats for duck and geese were collected and composited into 10 samples for duck and 10 samples for geese. Composite samples of fecal slurries were collected from swine facilities, one inside the watershed and one outside (2 duplicate samples) and dairy cattle facilities (one inside the watershed and two outside (2 duplicate samples each) human residential septic cleanout trucks (3 samples) and influent of three separate municipal wastewater treatment plants (3 samples). A total of 20 g of each fecal sample other than litter from each site was collected and was placed in a 20 ml, sterile,

polystyrene tube containing 10 ml of 20% glycerol and shipped on dry ice to the laboratory. All fecal samples were homogenized in the glycerol before DNA extraction. Discrete water samples from larger rivers and lakes were collected using a Van Dorn water sampler or with a churn splitter for discrete or composite samples. Samples from larger rivers were typically composites of 3 samples collected on a transect across the width of the river channel. Samples from smaller rivers were collected using automated samplers. Samples collected during high flow events were composited based on flow volume. Base flow samples were collected as grab samples. River samples were placed into sterile 1-L polystyrene bottles in duplicate and shipped on ice to the laboratory where they were filtered. Runoff samples from the litter application areas (e.g. edge of field runoff samples) were collected during or as soon as possible after rainfall events. Samples were collected either with a passive runoff collector for composite samples or with a dip sampler for discrete samples. Runoff samples were placed into sterile 1-L polystyrene bottles in duplicate and shipped on ice to the laboratory where they were filtered. Groundwater samples were collected directly from existing homeowner's wells or from hydraulically driven shallow probes. Spring samples were collected as grab samples or by using a peristaltic pump. All samples were placed into sterile 1-L polystyrene bottles and shipped on ice to the laboratory where they were filtered.

**Enumeration of Indicator Bacteria.** Indicator bacteria (fecal coliforms, *E. coli* and enterococci) were enumerated according to standard methods using multiple tube fermentation (MTF) and calculation of the most probable number according to SM-9221F or SM-9230 (APHA, 2005). MTF tubes containing *E. coli* were identified using broth cultures supplemented with (MUG) (SM-9221F) (2).



**Soil, Litter and Fecal Sample DNA Extraction.** Genomic DNA was extracted from soil, litter and fecal samples with Bio101 Fast®Spin® DNA extraction kits (QBiogene, Inc.) following the manufacturer's instructions. Typically 0.25 g of soil or litter was used in each extraction. DNA was purified by size-exclusion chromatography. Sepharose CL-4B (Sigma-Aldrich) was resuspended in Tris-HCL and sterilized by autoclave at 121 °C for at least 20 minutes. Micro-bio spin columns (Bio-Rad Laboratories) were packed with 1 mL of Sepharose CL-4B through centrifugation. Sepharose columns were then washed twice with Tris-HCl buffer (pH 8) and 50 to 150 µl of sample was added. Purified DNA was concentrated with ethanol precipitation and re-eluted in 100 µL sterile water.

**Water Sample DNA Extraction.** Within 12 hours of receipt at the laboratory all water samples were filtered through a sterile Supor-200, 0.2 µM filter and frozen at -80°C. Filters were then shattered with sterile glass beads and vortexed vigorously for 15 minutes with sterile, DNase, and RNase free water to remove solids and cells from the filters. The cell suspension was removed from the centrifuge tubes by pipette and placed in a 2 mL bead beating tube from the Bio101 Fast®Spin® DNA extraction kits. The cells were centrifuged at 20,000 x g for 10 minutes, and the supernatant was decanted. Genomic DNA was then extracted using the Bio101 Fast®Spin® DNA extraction kits (QBiogene, Inc). The extracted DNA was quantified using a Nanodrop® UV-Vis Spectrophotometer.

**T-RFLP Analysis.** Extracted genomic DNA and/or cloned DNA was amplified with phosphoramidite fluorochrome 5-carboxyfluorescein (FAM) labeled universal bacterial primers 8F-907R (16, 24), with *E.coli* genus specific primers (Tsen, et al. 1998), and *Bacteroidales* specific primers (Bernhard and Field, 2000). All PCR primers targeted the 16S rRNA gene. Triplicate PCR reactions were generated from each DNA extraction, combined and purified

145 using QIAquick PCR purification Kits (Qiagen). Approximately 200 ng each of PCR product  
146 was digested at 37°C for 6 hours with the *MspI* restriction enzyme (20µ/µL) (New England  
147 BioLabs). Samples were denatured by heating to 95° C for 3 minutes followed by cooling to  
148 4°C. The digested fragments were purified by ethanol precipitation.

149 **Primer Design.** Primers were designed using the ABI Primer Express v.2 program (Applied  
150 Biosystems, Foster City, CA) and were targeted to variable regions between the potential  
151 biomarker sequences and sequences of the top 20 closest related organisms in the GenBank  
152 database. The BLAST search (Basic Alignment Search Tool,  
153 <http://www.ncbi.nlm.nih.gov/blast/Blast.cgi>) was used to check the specificity of each primer.

154 **PCR Assay Conditions.** PCR was used to amplify approximately 900 bp of the 16S rRNA genes  
155 from *Bacteria* for clone library construction. Each 25 µL PCR reaction included 0.4 mg mL<sup>-1</sup>  
156 molecular-grade bovine serum albumin (BSA) (Sigma Chemicals), 1X PCR Buffer (Promega),  
157 1.5 mM MgCl<sub>2</sub>, 0.5 µM of both the forward (8F) (16) and reverse (907R) (24) primer  
158 (Invitrogen), 1U Taq DNA polymerase (Promega), 0.2 mM dNTP (Invitrogen), 1 µL DNA  
159 template, and molecular-grade water (Promega). Amplification was performed on a PerkinElmer  
160 Model 9600 thermocycler using the following conditions: 94 °C for 5 minutes, 30 cycles of 94  
161 °C (1 minute), 55 °C (45 seconds), and 72 °C (2 minute). A final extension at 72 °C for 7  
162 minutes was performed and the PCR products were held at 4°C. Specificity of the PCR primers  
163 to the poultry litter biomarker was evaluated with nested PCR by first amplifying non-target  
164 fecal samples by universal bacterial primers 8F, 907R and then amplifying by the potential  
165 poultry litter biomarker PCR primers. The nested PCR master mix and thermocycler conditions  
166 were similar to the universal PCR with the following exceptions: 1) forward and reverse PCR

primers were specific to the potential poultry biomarker as shown in Table 2, 2) the annealing temperature was 60 °C. Amplification by nested PCR was evaluated by gel electrophoresis.

**Clone Libraries.** Clone libraries were constructed from the original genomic DNA extracted from the soil and litter samples and amplified with either universal bacterial primers 8F-907R (16, 24), targeting the 16S rRNA genes of *Bacteria* or the *E. coli* genus specific primers V1SF-V3AR (41). The TOPO ® Cloning Reaction methods from Invitrogen™ were followed for clone library construction. Two clone libraries were constructed (targeting *Bacteria* and *E. coli*) from pooled DNA samples (i.e., 1 µl of genomic DNA extract from each sample was added to the PCR reaction for inclusion into the clones) based on the abundance of the various potential biomarkers as evidenced by the T-RFLP profiles.

**qPCR Assay Conditions.** Quantitative PCR (qPCR) was used to amplify 530 bp of the 16S rRNA gene from *Brevibacterium spp.* DNA samples were diluted to final concentrations of 3 ng/µL DNA. Each 25µL qPCR reaction included: 1X SYBR Green Master Mix (Roche), 0.5 µM of both the forward (LA35F) and reverse primer (LA35R) (Invitrogen), 5 % DMSO, 5 µL of diluted sample DNA, and molecular-grade water (Promega). Amplification was performed in triplicate on a Biorad Chromo4 thermocycler using the following conditions: 50 °C for 2 minutes, 95 °C for 15 minutes, 45 cycles of 95 °C (30 seconds), 60 °C (30 seconds), and 72 °C (30 seconds) with a plate read. The 45 cycles was followed by a final extension at 50 °C for 5 minutes. Immediately following the final extension was a melting curve from 70 °C to 90 °C, by 0.1 degree increments, holding for 5 seconds with a plate read. DNA standards ranging from  $6 \times 10^{-15}$  to  $10^{-21}$  ng/ul were prepared from serial dilutions of clone plasmid DNA containing the sequence of interest and used to develop the standard curve and method detection limit. Gene copy numbers were calculated from concentrations of positive control standards assuming 9.124

\*  $10^{14}$  bp/ul of DNA and one gene copy per genome. Detection limits for the qPCR assay were approximately 2000 plasmid copies in *E. coli*/L water and  $7.3 * 10^4$  plasmid copies in *E. coli*/gram of soil. Nested qPCR was performed by first amplifying DNA with the universal bacterial 16S rRNA 8F (16) and 907R (24) primers. The production of PCR products was confirmed on a 1.5% agarose gel. The 16S rRNA PCR products were purified with the QIAquick PCR purification kit (QIAGEN) were subjected to qPCR as previously described using the LA35F and LA35R primers for the poultry litter biomarker.

**Phylogeny.** The phylogeny of the LA35 clone was investigated using the following methods. The clone sequences were assembled and aligned with BioEdit v. 7.0.5.3 and sequences were checked for chimeras with the Ribosomal Database Project II Chimera Check program and Bellerophon. The 16S rRNA sequences of the closest neighbors to the clone sequences were downloaded for inclusion in the phylogenetic analysis. Multiple sequence alignments were constructed with Clustal W alignment tool and manually aligned in BioEdit. The bootstraps (1000 resamplings), maximum likelihood and distance matrix analysis (Kimura), and the reconstruction of the phylogenetic trees (FITCH) were performed with the Phylip 3.65 package and in particular the programs SEQBOOT, DNAML, DNADIST, FITCH, CONSENSE, and RETREE. The reconstructed phylogenetic tree was visualized with PhyloDraw V. 0.8 (Graphics Application Lab, Pusan National University).

## RESULTS

**Identification of potential biomarkers by T-RFLP.** A total of 20 T-RFLP profiles were generated from the 5 subsamples of each of the two litter and two soil samples. The T-RFs common among the subsamples and representing more than 1% of the community were selected

for cloning and sequencing (Table 1). A total of 3 *E. coli* T-RFs (i.e., T-RF 496.0, 498.9 and 500.8) and 3 *Bacteria* T-RFs (i.e., T-RF142.9, 147.3 and 158.9) were selected for cloning and sequencing. Clone libraries were constructed from PCR products amplified with *E. coli* specific primers (V1SF-V3AR) (41) or universal bacterial primers (8F-907R) (16, 24). A total of 300 plasmids from the clone libraries were randomly picked. T-RFLP analysis was carried out on each plasmid insert to identify which plasmids contained the T-RFs of potential biomarkers. Inserts containing the T-RFs of interest were sequenced and PCR primers were developed for those sequences containing mismatches as compared to BLAST database results of the top 20 closely related organisms. In all 4 PCR primers for members of 4 genera were developed; a *Brevibacterium* spp., a *Rhodoplanes* spp., a *Kineococcus* spp. and a *Pantoea ananatis* strain (Table 2). Two *E. coli* T-RFs were from plasmids that did not contain mismatches between the sequence of interest and the sequences of closely related organisms identified in a BLAST search and therefore were not appropriate biomarkers.

**Evaluation of biomarkers against fecal samples.** The PCR assays developed for the 4 potential biomarkers of poultry litter were tested for amplification against a variety of nontarget fecal samples from within and outside the watershed (Table 3). Only the *Brevibacterium* clone LA35 appeared to be a potential candidate biomarker for poultry litter in that did not amplify in any fecal samples with the exception of weak amplification in one duck and one goose sample from outside the watershed when analyzed with a nested PCR approach (i.e. PCR with universal bacterial primers and then with the *Brevibacterium* clone LA35 primers). The reconstructed phylogenetic tree of the *Brevibacterium* clone LA35 in relationship to other *Brevibacterium* spp. is presented in Figure 1.

**Quantification of the poultry litter biomarker in environmental samples.** A SYBR green qPCR protocol was developed and optimized using the LA35F and LA35R primers (Table 2) specific to the *Brevibacterium* clone LA35 poultry litter biomarker. The standard curve of the qPCR assay for the biomarker is presented in Figure 2. The detection limit of the qPCR assay was 6 gene copies/ul of extracted DNA.

Environmental samples from the potential poultry litter impacted watershed were tested for the presence of the biomarker with the qPCR assay (Table 4). A variety of samples from within the watershed were tested, some of which were expected to contain the biomarker (e.g., litter, contaminated soil, runoff samples), some of which had variable potential for higher biomarker levels (e.g., surface water), and some of which had lower potential for biomarker presence (i.e., groundwater samples).

The correlation between the poultry litter biomarker concentration (i.e., as quantified by qPCR) in water and litter samples and *E. coli* and *Enterococcus* as measured by most probable number is presented in Figures 3 and 4. In general the *Enterococcus* MPN counts were well correlated with the concentration of the biomarker in litter ( $R^2 = 0.75$ ) and with the biomarker concentration in water samples ( $R^2 = 0.89$ ). The correlation between *E. coli* concentrations and the biomarker in water samples was also strong ( $R^2 = 0.85$ ) while *E. coli* was less tightly (but significantly) correlated with the biomarker in litter samples ( $R^2 = 0.28$ ). Correlation of the biomarker with *E. coli* and *Enterococcus* spp. provides a line of evidence of the human health risk associated with the runoff from poultry litter application to fields although there is evidence that regrowth of these organisms is possible once they are introduced into the environment (36).

## DISCUSSION

The *Brevibacterium* sp. poultry litter biomarker developed in this study was validated in terms of sensitivity (100%) against numerous positive (poultry litter) samples from different locations with the watershed and for specificity (93.5%) against composite non-target fecal samples. These practices are in accordance with recent critical reviews (34, 40) that strongly recommend MST method validation. Future efforts will attempt to extend the method validation outside the watershed and possible outside the region as this biomarker could be useful for identifying fecal pollution sources in other river systems and coastal waters.

The *Brevibacterium* clone LA35 poultry litter biomarker was most closely related to *Brevibacterium avium*, which is associated with bumble-foot lesions in poultry (32). *Brevibacterium* spp. were recently identified in spent mushroom compost that was originally derived from chicken litter and cereal straw (29). Additionally *Brevibacterium avium*, *Brevibacterium iodinum*, and *Brevibacterium epidermidis* were found to represent more than 7% of a 16S rRNA clone library originating from broiler chicken litter (27). Certain *Brevibacterium* spp. are associated with milk and cheese curds(6), human skin(9), and soils (30). *Brevibacterium* spp. have been associated with disease in humans although to date these opportunistic pathogens have only been isolated from immunocompromised patients (4, 9, 18).

As poultry litter is land-applied as a disposal practice (19, 33, 35), it was important to identify a marker that could survive the process of deposition on bedding and spreading on fields.

Therefore, the T-RFLP screening process included both litter and contaminated soil samples.

This strategy allowed for the rapid elimination of numerous targets that could be abundant in the poultry fecal material, but not as abundant in the litter and not present in the environment after

litter application. This strategy for marker identification is in contrast with the work by Lu and colleagues (2007) where a genome fragment enrichment method was used to identify microbial sequences specific to chicken feces. Based on the PCR assays developed from clone libraries of the genome fragments, 6 to 40% of the chicken fecal samples collected from a wide geographic region contained DNA that could be amplified by the various assays (26). In comparison the LA35 biomarker was found in all the poultry litter samples tested, although it should be noted that all of the samples were collected in the Oklahoma/Arkansas region.

The examination of environmental samples from within the poultry litter impacted watershed suggest a correlation between the application of poultry litter to a field and concentration of the biomarker in the receiving waters, as evidenced by the generally decreasing trend in biomarker concentration with decreasing concentration of fecal indicator organisms. These results indicate that the watershed is in fact being impacted by the application of poultry litter to fields within the watershed. However, the magnitude of the impact as measured by the distribution of the biomarker within the watershed cannot be quantified with the limited number of environmental samples processed to date. Future work will include the testing of environmental samples from within the watershed by the qPCR assay to evaluate the distribution of the poultry litter-specific biomarker as compared to indicator bacteria, antibiotics and heavy metals. Additionally, testing of the poultry litter-specific biomarker against more fecal samples from other watersheds and additional avian fecal material will be conducted as the LA35 poultry litter biomarker was found in low abundance (i.e., a nested PCR approach was required for detection) in two non-target composite avian fecal samples (i.e., a duck and a goose sample) from outside the watershed.

## **Conclusions**



299 In summary a novel biomarker of poultry litter was identified and a 16S rRNA based real-time  
300 PCR assay was developed for this biomarker. The specificity of the assay (93.5%) was tested  
301 against 31 separate non-target fecal samples and sensitivity was tested against 10 target litter  
302 samples (100%). The field applicability of the assay was evaluated by testing for the biomarker  
303 in environmental samples expected to have variable concentrations of the biomarker, which we  
304 hypothesized would be correlated with the concentration of fecal indicator bacteria. A generally  
305 positive correlation was found between biomarker concentration and fecal indicator bacteria  
306 concentration which was particularly strong for enterococci. The research presented herein is the  
307 first identification of a *Brevibacterium* spp. for microbial source tracking studies and is among  
308 the first quantifiable method for tracking of poultry fecal sources in environmental waters.

#### 309 **ACKNOWLEDGMENTS**

310 This research was conducted in connection with work performed as retained experts in a pending  
311 legal case brought by the State of Oklahoma against several poultry integrators. Drs. Harwood  
312 & Olsen have been retained to serve as expert witnesses by the State of Oklahoma and have  
313 provided testimony regarding this research.

314 The authors are grateful for the assistance provided by Kyle Collins, William Blackmore, James  
315 Jackson, Erin O'Leary Jeapson and Michelle Andrews. Additionally the authors acknowledge  
316 the Molecular Research Core Facility at Idaho State University for graciously allowing us the use  
317 of their laboratory space and equipment.

## References

1. **Anderson, K., J. Whitlock, and V. Harwood.** 2005. Persistence and Differential Survival of Fecal Indicator Bacteria in Subtropical Waters and Sediments. *Applied and Environmental Microbiology* **71**:3041-3048.
2. **APHA.** 2005. Standard methods for the examination of water and wastewater, 21st ed. American Public Health Association, Inc., Washington, D.C.
3. **Bernhard, A., and K. Field.** 2000. Identification of Nonpoint Sources of Fecal Pollution in Coastal Waters by Using Host-Specific 16S Ribosomal DNA Genetic Markers from Fecal Anaerobes. *Applied and Environmental Microbiology* **66**:1587-1594.
4. **Beukinga, I., H. Rodriguez-Villalobos, A. Deplano, F. Jacobs, and M. Struelens.** 2005. Management of long-term catheter-related *Brevibacterium* bacteraemia. *Clinical Microbiology and Infection* **10**:465-467.
5. **Bitzer, C., and J. Sims.** 1988. Estimating the availability of nitrogen in poultry manure through laboratory and field studies. *Journal of Environmental Quality* **17**:47-54.
6. **Brennan, N., A. Ward, T. Beresford, P. Fox, M. Goodfellow, and T. Cogan.** 2002. Biodiversity of the bacterial flora on the surface of a smear cheese. *Applied and Environmental Microbiology* **68**:820-830.
7. **Bujozcek, G., J. Oleszkiewicz, R. Sparling, and S. Cenkowski.** 2000. High Solid Anaerobic Digestion of Chicken Manure. *Journal of Agricultural Engineering Research* **76**:51-60.
8. **Call, D., D. Satterwhite, and M. Soule.** 2007. Using DNA suspension arrays to identify library-independent markers for bacterial source tracking. *Applied and Environmental Microbiology* **41**:3740-3746.

- 341 9. **Cannon, J., F. Spadoni, S. Pesh-Iman, and S. Johnson.** 2005. Pericardial infection  
342 caused by *Brevibacterium casei*. *Clinical Microbiology and Infection* **11**:164-165.
- 343 10. **Carson, C., B. Shear, M. Ellersieck, and A. Asfaw.** 2001. Identification of Fecal  
344 *Escherichia coli* from Humans and Animals by Ribotyping. *Applied and Environmental*  
345 *Microbiology* **67**:1503-1507.
- 346 11. **Connor, R., M. Connor, K. Irgolic, J. Sabrsula, H. Gurleyuk, R. Brunette, C.**  
347 **Howard, J. Garcia, J. Brien, J. Brien, and J. Brien.** 2005. Transformations, Air  
348 Transport and Human Impact of Arsenic from Poultry Litter. *Environmental Forensics*  
349 **6**:83-89.
- 350 12. **Desmarais, T., H. Solo-Gabriele, and C. Palmer.** 2002. Influence of Soil on Fecal  
351 Indicator Organisms in a Tidally Influenced Subtropical Environment. *Applied and*  
352 *Environmental Microbiology* **68**:1165-1172.
- 353 13. **Dick, L. K., A. E. Bernhard, T. J. Brodeur, J. W. Santo Domingo, J. M. Simpson, S.**  
354 **P. Walters, and K. G. Field.** 2005. Host distributions of uncultivated fecal *Bacteroidales*  
355 bacteria reveal genetic markers for fecal source identification. *Applied and*  
356 *Environmental Microbiology* **71**:3184-3191.
- 357 14. **Dombek, P., L. Johnson, S. Zimmerley, and M. Sadowsky.** 2000. Use of repetitive  
358 DNA sequences and the PCR to differentiate *Escherichia coli* isolates from human and  
359 animal sources. *Applied and Environmental Microbiology* **66**:2572-2577.
- 360 15. **Edwards, D., and T. Daniel.** 1994. A comparison of runoff quality effects of organic  
361 and inorganic fertilizers applied to fescue grass plots. *Journal of the American Water*  
362 *Resources Association* **30**:35-41.

16. **Edwards, U., T. Rogall, H. Blocker, M. Emde, and E. Bottger.** 1989. Isolation and  
direct complete nucleotide determination of entire genes. Characterization of a gene  
coding for 16S ribosomal RNA. *Nucleic Acids Research* **17**:7843-7853.
17. **Field, K., and M. Samadpour.** 2007. Fecal source tracking, the indicator paradigm, and  
managing water quality. *Water Research* **41**:3517-3538.
18. **Janda, W., P. Tipirneni, and R. Novak.** 2003. *Brevibacterium casei* Bacteremia and  
Line Sepsis in a Patient with AIDS. *Journal of Infection* **46**:61-64.
19. **Jenkins, M., D. Endale, H. Schomber, and R. Sharpe.** 2006. Fecal bacteria and sex  
hormones in soil and runoff from cropped watersheds amended with poultry litter.  
*Science of the Total Environment* **358**:164-177.
20. **Kelleher, B., J. Leahy, A. Henihan, T. O'Dwyer, D. Sutton, and M. Leahy.** 2002.  
Advances in poultry litter disposal technology – a review. *Bioresource Technology*  
**83**:27-36.
21. **Kelley, T., O. Pancorbo, W. Mercka, S. Thompson, M. Cabrera, and H. Barnhart.**  
1994. Fate of Selected Bacterial Pathogens and Indicators in Fractionated Poultry Litter  
During Storage. *Journal of Applied Poultry Research* **3**:279-288.
22. **Kildare, B. J., C. M. Leutenegger, B. S. McSwain, D. G. Bambic, V. B. Rajal, and S.  
Wuertz.** 2007. 16S rRNA-based assays for quantitative detection of universal, human-,  
cow-, and dog-specific fecal *Bacteroidales*: a Bayesian approach. *Water Research*  
**41**:3701-3715.
23. **Kuntz, R., P. Hartel, J. Rodgers, and W. Segars.** 2004. Presence of *Enterococcus*  
*faecalis* in broiler litter and wild bird feces for bacterial source tracking. *Water Research*  
**38**:3551-3557.

- 386 24. **Lane, D.** 1991. 16S/23S rRNA sequencing. *In* E. Stackebrandt and M. Goodfellow (ed.),  
 387 Nucleic acid sequencing techniques in bacterial systematics. John Wiley and Sons, New  
 388 York, N.Y.
- 389 25. **Leclerc, H., D. Mossel, S. Edberg, and C. Struijk.** 2001. Advances in the Bacteriology  
 390 of the Coliform Group: Their Suitability as Markers of Microbial Water Safety. *Annual*  
 391 *Reviews in Microbiology* **55**:201-234.
- 392 26. **Lu, J., J. Domingo, and O. Shanks.** 2007. Identification of a chicken-specific fecal  
 393 microbial sequences using a metagenomic approach. *Water Research* **41**:3561-3574.
- 394 27. **Lu, J., S. Sanchez, C. Hofacre, J. Maurer, B. Harmon, and M. Lee.** 2003. Evaluation  
 395 of Broiler Litter with Reference to the Microbial Composition as Assessed by Using 16S  
 396 rRNA and Functional Gene Markers. *Applied and Environmental Microbiology* **69**:901-  
 397 908.
- 398 28. **Mozaffari, M., and J. Sims.** 1994. Phosphorus availability and sorption in an Atlantic  
 399 coastal plain watershed dominated by animal-based agriculture. *Soil Science* **157**:97-107.
- 400 29. **Ntougias, S., G. Zervakis, N. Kavroulakis, C. Ehaliotis, and K. Papadopoulou.** 2004.  
 401 Bacterial Diversity in Spent Mushroom Compost Assessed by Amplified rDNA  
 402 Restriction Analysis and Sequencing of Cultivated Isolates. *Systematic and Applied*  
 403 *Microbiology* **27**:746-754.
- 404 30. **Onraedt, A., W. Soetaert, and E. Vandamme.** 2005. Industrial importance of the genus  
 405 *Brevibacterium*. *Biotechnology Letters* **27**:527-533.
- 406 31. **Parveen, S., R. Murphree, L. Edmiston, C. Kaspar, and M. Tamplin.** 1999.  
 407 Discriminant analysis of ribotype profiles of *Escherichia coli* for differentiating human

and nonhuman sources of fecal pollution. *Applied and Environmental Microbiology*  
**65**:3142-3147.

32. **Pascual, C., and M. Collins.** 1999. *Brevibacterium avium* sp. nov., isolated from poultry. *International Journal of Systematic Bacteriology* **49**:1527-1530.
33. **Pirani, A., K. Brye, T. Daniel, B. Haggard, E. Gbur, and J. Mattice.** 2006. Soluble Metal Leaching from a Poultry Litter–Amended Udult under Pasture Vegetation. *Vadose Zone Journal* **5**:1017-1034.
34. **Santo Domingo, J., D. Bambic, T. Edge, and S. Wuertz.** 2007. Quo vadis source tracking? Towards a strategic framework for environmental monitoring of fecal pollution. *Water Research* **41**:3539-3552.
35. **Schroeder, P., D. Radcliffer, and M. Cabrera.** 2004. Rainfall Timing and Poultry Litter Application Rate Effects on Phosphorus Loss in Surface Runoff. *Journal of Environmental Quality* **33**:2201-2209.
36. **Scott, T., J. Rose, T. Jenkins, S. Farrah, and J. Lukasik.** 2002. Microbial Source Tracking: Current Methodology and Future Directions. *Applied and Environmental Microbiology* **68**:5796-5803.
37. **Scott, T. M., T. M. Jenkins, J. Lukasik, and J. B. Rose.** 2005. Potential use of a host associated molecular marker in *Enterococcus faecium* as an index of human fecal pollution. *Environmental Science and Technology* **39**:283-287.
38. **Seurinck, S., T. Defoirdt, W. Verstraete, and S. D. Siciliano.** 2005. Detection and quantification of the human-specific HF183 *Bacteroides* 16S rRNA genetic marker with real-time PCR for assessment of human faecal pollution in freshwater. *Environmental Microbiology* **7**:249-259.

- 431 39. **Sharpley, A., T. Daniel, J. Sims, and D. Pote.** 1996. Determining environmentally  
432 sound soil phosphorus levels. *Journal of Soil and Water Conservation* **51**:160-168.
- 433 40. **Stoeckel, D., and V. Harwood.** 2007. Performance, design and analysis in microbial  
434 source tracking studies. *Applied and Environmental Microbiology* **73**:2405-2415.
- 435 41. **Tsen, H., C. Lin, and W. Chi.** 1998. Development and use of 16S rRNA gene targeted  
436 PCR primers for the identification of *Escherichia coli* cells in water. *Journal of Applied*  
437 *Microbiology* **85**:554-560.
- 438 42. **USEPA.** 2000. Improved enumeration methods for the recreational water quality  
439 indicators: enterococci and *Escherichia coli*. EPA-821/R-771 97/004. U.S.  
440 Environmental Protection Agency.
- 441 43. **USEPA.** 2005. Microbial source tracking guide document, EPA/600/R-05/064. U.S.  
442 Environmental Protection Agency.
- 443 44. **USEPA.** 2001. Protocol for developing pathogen TMDLs. EPA 841-R-00-002. U.S.  
444 Environmental Protection Agency.
- 445 45. **Van Donsel, D., E. Geldreich, and N. Clarke.** 1967. Seasonal Variations in Survival of  
446 Indicator Bacteria in Soil and their Contribution to Storm-water Pollution. *Applied*  
447 *Microbiology* **15**:1362-1370.
- 448 46. **Wade, T., R. Calderon, E. Sams, M. Beach, K. Brenner, A. Williams, and A. Dufour.**  
449 2006. Rapidly measured indicators of recreational water quality are predictive of  
450 swimming-associated gastrointestinal illness. *Environ. Health Perspectives* **114**:24-28.
- 451 47. **Wuertz, S., and J. Field.** 2007. Emerging microbial and chemical source tracking  
452 techniques to identify origins of fecal contamination in waterways. *Water Research*  
453 **41**:3515-3516.

Table 1. Common T-RFs among replicates from two fecal-contaminated poultry litter samples  
and two soils to which the litter had been applied.

T-RF	Number of subsamples tested (number containing T-RF of interest)			
	Litter A	Litter B	Soil A	Soil B
<i>E.coli</i> PCR products, digested with <i>MspI</i>				
<u>496.0</u>	4 (4)	5 (4)	5 (3)	5 (5)
<u>498.9</u>	4 (4)	5 (5)	5 (4)	5 (5)
<u>500.8</u>	4 (4)	5 (5)	5 (5)	5 (5)
Universal bacteria PCR products, digested with <i>MspI</i>				
80.1	4 (4)	5 (5)	5 (0)	3 (3)
130.9	4 (3)	5 (5)	5 (1)	3 (0)
<u>142.9</u>	4 (4)	5 (4)	5 (2)	3 (2)
<u>147.3</u>	4 (4)	5 (5)	5 (5)	3 (2)
<u>158.9</u>	4 (3)	5 (5)	5 (4)	3 (2)
165.0	4 (3)	5 (5)	5 (4)	3 (2)
*Underlined T-RFs correlate to those organisms for which PCR primers were developed				



459 Table 2. Nucleotide sequences and targets of primers used in this study.

460

Primer	Target	Sequence (5'-3')	Position	T <sub>m</sub> (°C)	T-RF
LA35F	<i>Brevibacterium</i>	ACCGGATACGACCATCTGC	166-184	57	147.3
LA35R	clone LA35	TCCCCAGTGTCAGTCACAGC	717-736	58	
SA19F	<i>Kineococcus</i>	TACGACTCACCTCGGCATC	163-181	56	158.9
SA19R	<i>spp.</i>	ACTCTAGTGTGCCCCGTACCC	602-621	55	
SB37F	<i>Rhodoplanes</i>	AACGTGCCTTTTGGTTTCG	143-160	56	142.9
SB37R	<i>spp.</i>	GCTCCTCAGTATCAAAGGCAG	616-626	55	
SA15F	<i>Pantoea</i>	CGATGTGGTTAATAACCGCAT	490-510	56	500.8
SA15R	<i>ananatis</i>	AAGCCTGCCAGTTTCAAATAC	668-688	55	

461

462 Table 3. Specificity of the poultry litter biomarker assay tested against fecal samples from within and outside the watershed.

463

Fecal sample (inside or outside watershed)	Number of samples tested (Number of samples containing potential biomarker)			
	<i>Brevibacterium</i> clone	<i>Rhodoplanes</i> clone	<i>Kineococcus</i>	<i>Pantoea ananatis</i>
	LA35	SB37	clone SA19	clone SA15
Beef cattle (outside)	5 (0)	5 (2)	5 (1)	5 (0)
Beef cattle (inside)	5 (0)	5 (3)	5 (5)	5 (1)
Dairy cattle (outside)	2 (0)	2 (1)	2 (1)	2 (1)
Dairy cattle (inside)	1 (0)	1 (1)	1 (0)	1 (0)
Swine (outside)	1 (0)	1 (1)	1 (1)	1 (0)
Swine (inside)	1 (0)	1 (0)	1 (0)	1 (0)
Duck (outside)	2 (1)*	2 (2)	2 (2)	2 (2)
Duck (inside)	3 (0)	3 (1)	3 (1)	3 (2)
Goose (outside)	3 (1)*	3 (3)	3 (2)	3 (2)
Goose (inside)	2 (0)	2 (2)	2 (1)	2 (1)
Human sewage (outside)	2 (0)	2 (2)	2 (2)	2 (1)
Human sewage (inside)	4 (0)	4 (3)	4 (1)	4 (1)
* One duplicate amplified when analyzed with a nested PCR assay.				





Figure 1. Reconstructed phylogenetic tree of the *Brevibacterium* spp. based on 16S rRNA.

Numbers at the nodes represent bootstrap values (i.e. the number of times this organism was found in this position relative to other organisms in 1000 resamplings of the data). Bootstraps less than 50% are not shown. The closest cultured organisms as reported in an NCBI BLAST search are reported. The distance bar represents a 1% estimated sequence divergence.

473



474

475

476

477 Figure 2. Standard curve of measured Ct values and standard deviations versus log plasmid

478 biomarker concentration.


Table 4. Environmental samples tested for *Brevibacterium* clone LA35 poultry litter biomarker

Sample type	Number	% of samples		Range of biomarker present (16S rRNA copies/L water or g soil or g litter)
	samples tested	containing biomarker <sup>a</sup>	% of samples quantifiable <sup>b</sup>	
Litter	10	100	100	$2.2 \times 10^7 \pm 7.1 \times 10^6 - 2.5 \times 10^9 \pm 9.5 \times 10^7$
Soil	10	100	50	$7.0 \times 10^3 \pm 4.4 \times 10^2 - 2.9 \times 10^5 \pm 2.0 \times 10^4$
Edge of field runoff	10	100	100	$2.6 \times 10^3 \pm 1.2 \times 10^2 - 5.5 \times 10^7 \pm 5.3 \times 10^6$
River	10	50	20	$2.9 \times 10^3 \pm 8.6 \times 10^2 - 3.2 \times 10^4 \pm 6.8 \times 10^3$
Groundwater	6	0	0	Not applicable

<sup>a</sup> indicates the percent of samples in which the biomarker was identified by qPCR or nested qPCR methods

<sup>b</sup> indicates the percent of samples for which a quantifiable number of biomarker genes were measured by qPCR

482



483

484 Figure 3. Correlation between the concentrations of poultry litter biomarker, *E. coli* and485 *Enterococcus* spp. in poultry litter samples.




Figure 4. Correlation between the concentrations of poultry litter biomarker, *E. coli* and *Enterococcus* spp. in water samples.